




MANAGEMENT GUIDE

**M77 MALE MATED WITH
CONVENTIONAL FEMALES
PARENT STOCK**



1.	KEY MESSAGES FOR REARING PERIOD (0-22 WEEKS): EQUIPMENT AND STOCKING DENSITY	3
2.	KEY MESSAGES FOR THE PRODUCTION PERIOD (22-64 WEEKS): EQUIPMENT AND STOCKING DENSITY	4
3.	REARING PERIOD (0-22 WEEKS)	5
	HOW TO HAVE A SUCCESSFUL BROODING PERIOD?	5
	HOW TO FOLLOW THE BODYWEIGHT TARGETS?	6
	HOW TO REACH AND MAINTAIN A GOOD FLOCK UNIFORMITY?	7
	HOW TO PROMOTE A GOOD BEHAVIOUR AND PROVIDE A GOOD ANIMAL WELFARE?	8
4.	PRODUCTION PERIOD (22-64 WEEKS)	9
	HOW TO HAVE A SUCCESSFUL MIXING PERIOD AND GOOD EARLY FERTILITY?	9
	HOW TO MAXIMIZE THE PERSISTENCY OF FERTILITY?	10
5.	BIOSECURITY	11

1. KEY MESSAGES FOR REARING PERIOD (0-22 WEEKS): EQUIPMENT AND STOCKING DENSITY



- Successful rearing = More than 80% males with a good sexual development at the time of the light stimulation (154-161 days of age) and in line with female maturity. All males should show straight legs, correct toes, beak, body conformation and feathering.
- Flock management to be adapted according to the local rearing conditions (house type, climate...).
- 1 Ad libitum feeding during the first 2 weeks and until 3 weeks of age if the 2 weeks bodyweight target is not achieved. Review again at 21 days before controlling feed amounts.
 - 2 Combine the reduction of day length and light intensity in the first 2 weeks: 24 h light/60 lux → 8 h light/5-10 lux. If the males and females are brooded in the same house, then use the same lighting program as for the Efficiency Plus females. In separate houses, it may be possible to delay the step down of light if growth is below target.
 - 3 Provide 3-5 cm / male of perch space or 1-2 m² of platform / 500 males from 28 days to train the males to jump up to the slats.
 - 4 Grade or at least isolate the smallest chicks from 7 days. Four weeks after the 4th week grading, the pen with the small males must reach the 8 weeks bodyweight target to secure skeletal development which is almost completed at this age.
 - 5 Maintain 45-60 minutes eating time; adjust the feeding programme to achieve this – (subject to local regulations).
 - 6 Avoid disruption in growth between 16-22 weeks (+150-160 g week), especially when changing feed type and after vaccinations. Assess body conformation on a regular basis. All the life of the flock:

- 0-4 weeks:** Obtain a good start and a successful grading
- 5-10 weeks:** Develop the skeleton of the birds
- 11-15 weeks:** Control the muscle development
- 16-22 weeks:** Prepare the males for a proper development of the testicles

EQUIPMENT AND STOCKING DENSITY DURING REARING	Stocking density *	3-4 males/m ²
	Brooders	1 for 500 chicks
	Watering - round	1 for 80 birds
	Watering - nipple	1 for 8-10 birds
	Feeding - chain	20 cm/male (10 m/100 males)
	Feeding pans - round	1 for 8 males
	Feeding pans - oval	1 for 8-10 males
	Spin feeder	1 for 1000-1200 males
	Feed distribution time	< 4 minutes

* Stocking density may be subject to local regulations which must be observed at all times.



i



[Management guide Efficiency Plus PS](#)



[Poster Feed distribution](#)



[Performance Objectives M77 PS male mated with conventional females](#)



[Technical bulletin Grading](#)

- ▷ Assure good distribution of both feed and water for a good flock uniformity.
- ▷ Where possible males should be grown using the same type of feeder in rearing and in production.
- ▷ Maintain good litter quality.
- ▷ Meet the environmental and air quality requirements for a healthy flock.
- ▷ Weigh 3-5% of the flock and at least 60 birds per pen per week.

2. KEY MESSAGES FOR THE PRODUCTION PERIOD (22-64 WEEKS): EQUIPMENT AND STOCKING DENSITY

KEY POINTS

EQUIPMENT AND STOCKING DENSITY DURING PRODUCTION	Watering - round	1 for 80 males
	Watering - nipple flow 70 - 100 ml/min*	1 for 6 to 10 males
	Feeding - chain	20 cm feeder space per male / 10 m length for 100 males
	Feeding pans - round ø 35 cm - 13.8 in	1 for 8 males
	Feeding pans - oval	1 for 8-10 males
	Feed distribution time	< 4 minutes
	Light intensity	60-80 lux

* Some nipple drinker systems are designed to operate with lower flow rates for breeders so check the manufacturer's recommendations or seek advice from your Hubbard Technical Manager.

LIGHT

► Same procedures as for females. To assure a good balance of the male and female sexual maturity, males may sometimes be stimulated one week earlier than females.

TRANSFER TIME (20-22 WEEKS) TO 27 WEEKS OF AGE

► Avoid disruption in growth between 20-27 weeks, especially when changing feed type (Male 1 diet from 20 weeks of age) and mixing males. Ensure the weekly growth rate is on target.

► Mixing is crucial to establish a good relationship between the males and females:

- ▷ Mix males that are on target bodyweight with uniform skeleton size and good shank length (10 weeks bodyweight achieved to maximise skeletal maturity). Never mix shy and/or immature males.
- ▷ Progressive mixing is preferred: 5% of males between 22-24 weeks and gradual increase to 8.5% at 25 weeks of age. More than 8.5% males could be a risk to maintain good female feathering and low floor eggs %.
- The risk of excessive weight gain and / or loss of condition up to 28 weeks is high:
 - ▷ Use a proper grill size 45 mm x 60 mm or adjust pan feeders to control male access to the female feeder. Also ensure that the corners of the feed track are sealed with covers. Check behaviour throughout the day to see if males are stealing feed.
 - ▷ Check feed distribution at least 2 times per week at first and then every week. Feed males after the female feed distribution is complete.
 - ▷ Weigh 3-5% of the flock and at least 60 males per pen weekly. Adjust feed according to both condition and bodyweight.

AFTER 28 WEEKS OF AGE

► Male bodyweight gain should be regular and within the range shown on the graph of the Performance Objectives. Feed to maintain their condition. → See Performance Objectives "M77 PS male mated with conventional females".

► Use the low protein male feed (Male 2 diet) which is one option to help to maintain male in good condition. → See "Conventional Nutrition Recommendations".


► Male replacement to maintain fertility after 40-45 weeks of age only if all biosecurity risk controls at all times are respected. → See Technical Bulletin "Male replacement".

▷ "Intra-farm" male replacement where males can be exchanged between houses.


▷ External flock replacement where the males should come from a known source with known health status.

WATER


► Control water quality: pH = 5.5 to 6.8/Oxidation Reduction Potential (ORP) or Redox potential > 600 mV with chlorine/Chlorine concentration at the end of the pipe = 1 ppm (Acceptable range: 0.5 to 3 ppm). → Maintain a good litter quality.




Growth targets for M77 during the production period




[Management guide Efficiency Plus PS](#)




[Conventional Nutrition Recommendations](#)



[Performance Objectives M77 PS male mated with conventional females](#)




[Technical bulletin Water Quality](#)



[Technical bulletin Male replacement](#)

3. REARING PERIOD (0-22 WEEKS): HOW TO HAVE A SUCCESSFUL BROODING PERIOD?

The key management practices are shown in our brooding poster. → See Poster *“Brooding PS”*. Close observation during the first 10 days of the chicks in their environment will make brooding successful by adjusting light, feed, water, equipment, heat and minimum ventilation.



Optimal crumble starter feed

KEY POINTS

- If possible, use an intermittent lighting programme to promote early feed intake (e.g. cycles of 2.30h light/30 min dark from day 1 to day 7) and always respect the recommended hours of light below and a minimum of 4 cycles of light dark per day (if permitted by local regulations).
- Chick paper on at least 50% of the surface with feed.
- Avoid a build-up of feed dust during the first 14 days.
- Use of all the floor space by 10 days if brooders are used.
- Ideal pH: 5.5 to 6.8 with an optimal ORP or Redox potential higher than 600 mV (with chlorination) or chlorine concentration at the end of the pipe = 1 ppm (acceptable range: 0.5 to 3 ppm). Water at ambient temperature: 24-26°C.
- Grade or at least isolate the smallest birds from 7 days.
- Keep using starter crumbles if bodyweight not on target at 28 days and use grower feed if bodyweight is on target.
- The rate of step down of the photoperiod may often be slower (e.g. achieve 8 h light not earlier than 21 days) in the following cases:
 - ▷ For open-sided housing, allow birds to feed in the cooler part of the day. Also use the experience from the previous flocks.
 - ▷ When males and females are grown in the same house.
 - ▷ In case the target weight of the males is not achieved at 7 days.



AGE	Days	0	1	2	3	4	5	6	7	8	9	10	11	12	13	14
TEMPERATURE	Under the heaters	35-36			32-34			28-30								
	Living area	28						27-28			25-26					
	Whole house heating	31-32	30-31	29-30	28-29	26-27										
HUMIDITY	%	50-60														
EQUIPMENT	Drinker	1 round drinker for 60 birds; 1 nipple for 8 birds									1 round drinker for 80 birds; 1 nipple for 8-10 birds					
	Feeder	1 feeder/50-70 birds + chick paper (> 50%)									Chain feeder: 6-8 cm/bird; 1 oval pan feeder/13-14 birds*; 1 round pan feeder/12 birds					
DENSITY	Chicks/m ²	25-30			12			6-8								
LIGHT PROGRAMME	Light duration (h)	24	22	21	20	19	18	17	16	15	14	13	12	11-12 **	8-12 **	
	Intensity *** (lux)	60						40			20		5-10			
FEED	Ad libitum	Ad libitum for 2 weeks and until 3 weeks if the 2 weeks bodyweight is not achieved. Review again at 21 days before controlling feed amounts.														
	Feed type	Pre-starter crumbles or Mini pellets (< 2.5 mm diameter) for a minimum of 14 days														

* Feeding space to be adjusted according to the behaviour of the birds.
 ** In case the target weights of the males are not achieved at 7 days.
 *** If permitted by local regulations.

3. REARING PERIOD (0-22 WEEKS): HOW TO FOLLOW THE BODYWEIGHT TARGETS?

MANUAL WEIGHING




Every week, always the same day: hatch day is the best.



Best before feeding.

After feeding, use a with feed and water bodyweight target.



- Check the calibration of the scale before weighing the birds.
- Take samples from 3 separate places but not too close to the main feed hopper. Always keep the same location.
- **Sample size: 3-5% of the flock size and at least 60 birds per pen.**
- Weigh all the birds of the catching pen.

AUTOMATIC WEIGHING




Every week, always the same day: hatch day is the best.



Best before feeding.

After feeding, use a with feed and water bodyweight target.



Results from an automatic scale - good bodyweight distribution

- Ensure to place the scales as soon as possible to maximize the number of birds weighed.
- Always check the number of birds weighed per pen (at least 300 birds/weighing) and make sure that the bodyweight distribution shows a “bell curve” shape (see figure above). The lower and upper limits of the bodyweights often need to be adjusted.
- If an unexpected variation of the bodyweight from the objective is observed, it is recommended to re-weigh by hand a new sample.
- Check weekly the calibration of the platform scales.



Analysis of the weekly results for an optimal feed allocation management:

1. In order to define properly the feed allocation per bird, the primary objective is to use the bodyweight average calculated once the weighing has been completed.
2. Compare the average bodyweight with the bodyweight target. Plot the average weight on the growth curve to see the trend.
3. Calculate the weekly gain and compare it with the objective mentioned in the Performance Objectives *“M77 PS male mated with conventional females”*.
4. Adjust feed allocation according to the weekly weight gain achieved and the weight gain targeted for the next week. Please do not strictly follow the feed intake targets as they are only provided as a guideline and consider actual feed increases.

3. REARING PERIOD (0-22 WEEKS): HOW TO REACH AND MAINTAIN A GOOD FLOCK UNIFORMITY?

There are two possible indicators to express flock uniformity:

1. The uniformity is defined by the percentage of birds weighing between +/- 10% around the average flock bodyweight. → See chart below.
2. The coefficient of Variation (CV) is the variation of bodyweights within the flock (= Standard deviation/average flock bodyweight). The lower the CV%, then the flock is more uniform.

GRADING

➤ 7 days: grade or at least isolate the smallest chicks by eye and manage them with special care to achieve target weight by 4 weeks.



➤ 21-28 days: grade 100% of the flock and create weight groups, each with at least 85% uniformity (CV = 7.5). → See Technical Bulletin "Grading".

➤ 29-84 days: aim to maintain or improve flock uniformity by having a close follow up of the feed distribution. → See Poster "Feed distribution". Poor wing feather quality may be an indication of poor feed distribution or feeder management.

➤ 84-98 days: if flock uniformity is less than 80% (CV > 8), re-grade the flock to assure that each weight group achieves at least 85% uniformity (CV < 7.5). Have a close bodyweight follow up during the vaccination periods. Anticipate this with extra feed.

FEED PROGRAMME

➤ Daily feeding gives successful results in many situations.

➤ Fractionated feeding (if permitted by local regulations) could be required to maintain a correct clearance time (45-60 min) and a proper eating behaviour. It will provide larger amount of feed during the days with feed.

- The 6/1 feeding programme means 6 days with feed and one day off.
- The 5/2 feeding programme means 5 days with feed and two days off.
- The alternate feeding strategy (SAD) means that birds are fed every other day.

➤ Example at 5 weeks: daily feeding with 50 g/bird/day equal to:

➤ 58,3 g/bird/feed day with a 6/1 feeding programme = (50 g x 7 days)/6 feed days.

➤ 70 g/bird/feed day with a 5/2 feeding programme = (50 g x 7 days)/5 feed days.

➤ Whichever the option chosen, it is important to use transitions as shown below and not exceed 150 g/feed day/bird.


➤ Option N° 1: daily → 6/1 (from 4 weeks of age) → 5/2 (from 5 weeks of age).

➤ Option N° 2: daily → 6/1 (from 4 weeks of age) → 5/2 (from 5 weeks of age) → 4/3 or SAD (from 6 weeks of age).


➤ Use of a low energy grower feed (< 2650 kcal/kg) is preferred to improve gut health and eating behaviour. If high fibre sources are added to the feed these must be of consistent quality and mycotoxin free. → See Technical Bulletin "Dietary fibre".




i




[Poster
Feed distribution](#)



[Technical bulletin
Grading](#)



[Technical bulletin
Dietary fibre](#)



Results of one manual weighing - % uniformity definition

3. REARING PERIOD (0-22 WEEKS): HOW TO PROMOTE A GOOD BEHAVIOUR AND PROVIDE A GOOD ANIMAL WELFARE?

For all enrichments it is important to consider the risks to compromise biosecurity, especially from straw or other bales and material brought into the house and also the ease of washing of fixed enrichments like platforms.


► PERCHES

From 28 days, the use of a perching system is strongly recommended to stimulate activity and to train birds jump onto the slats.

- ▷ Provide 3-5 cm of perch space per bird.
- ▷ Provide 1-2 m² of platform for 500 birds. Optimal position of the platform is under the drinking system once the birds are able to jump onto the slats.




- ▷ Platforms and perches should not have sharp edges that may damage birds.



► WATER MANAGEMENT / WATER QUALITY

- ▷ Provide *ad libitum* water during extremely hot weather or medication treatment.
- ▷ With a regular and proper management of drinker height and water flow according to flock age, no control access to water should be needed.
- ▷ Control access to water 2-3 hours after the end of the feed clean-up (if permitted by local regulations). The crop should be soft before closing the water.
- ▷ If using an alternate feeding strategy:
 1. On days with feed: control of the access to water is recommended.
 2. On days with no feed: give a minimum of 2 hours water and longer if required.



- ▷ Regularly check the chemical and bacteriological water quality to ensure that water sanitation is working properly:
 1. Optimal pH: 5.5 to 6.8.
 2. Optimal ORP or Redox potential higher than 600 mV with chlorine.
 3. Chlorine concentration at the end of the pipe: target = 1 ppm (acceptable range: 0.5 to 3 ppm).

► GRIT AND GRAIN - optional - consider the risks to biosecurity.


- ▷ Insoluble grit to promote the gizzard development:
 1. ϕ 2-3 mm: 3-5 g/bird/week from 5 weeks of age.
 2. ϕ 3-5 mm: 3-5 g/bird/week from 10 weeks of age.



- ▷ Scratch grain (cracked maize or whole wheat): 3 g/bird/day 4-5 h after feeding from 5 weeks of age.

► ENRICHMENT - optional unless required by local regulations.

- ▷ White ropes – max 20 cm long – no contact with the litter.
- ▷ Wood shaving balls – 1 piece for 500 to 1000 birds, placed on floor.
- ▷ Pecking blocks – 1 piece for 500 to 1000 birds – Consider the hardness of the material.
- ▷ Alfalfa balls – 1 piece for 500 to 1000 birds, placed on floor.



4. PRODUCTION PERIOD (22-64 WEEKS): HOW TO HAVE A SUCCESSFUL MIXING PERIOD AND GOOD EARLY FERTILITY?

OBJECTIVE: REACH MORE THAN 91.6% FERTILITY AT 27 WEEKS OF AGE.

➤ The objective is to avoid disruption in growth between 20-27 weeks which is a key period to develop the fertility potential of the males.

➤ A good broiler breeder male shows a good beak, long shanks, a good body condition (fleshing score: 2-3), good legs and feet (no bent toes, footpads with no injury), red comb and wattles, good feather cover with partial feather loss and a red and moist vent.

LIGHTING PROGRAM

➤ Use the lighting programme table of the female Efficiency Plus. → See *“Performance Objectives Efficiency Plus”*.

➤ If the sexual maturity is poor at 20-21 weeks of age (poor fleshing score: > 30% of the males with score 1), it is possible to give the first stimulation one week before the females.

MALE TRANSFER BEFORE THE FEMALES (20 TO 22 WEEKS)

➤ Where possible, it is a good practice to place males in the production house 3 to 5 days before the females to allow males to have a better adaptation to the new feeding and drinking systems (especially if spin feeding is used in rearing).

➤ Place males all along the house for a uniform distribution and not near a door.

➤ During the first days after the transfer, check if they find drinkers and feeders especially if changing compared to the rearing house. The male feeding system could be lowered to help them find the feed.

➤ Use Male 1 diet from 20 weeks of age.

MALE MIXING

➤ Mix males that are on target bodyweight with uniform skeleton size and good shank length. Never mix shy, immature or underweight males as they will be dominated by the females.

➤ For a better integration mix males ideally in the evening. Carefully observe the male / female interaction.

➤ Progressive mixing is ideal (2 or 3 steps):

▷ 5% between 22-24 weeks and gradual increase to 8.5% at 25 weeks of age. A place will be needed to store the spare males following proper conditions (density and equipment).

▷ More than 8.5% could be risky to maintain good female feathering and low floor eggs %.



Good broiler breeder male



[Performance Objectives M77 PS male mated with conventional females](#)

The risk of excessive weight gain and/or loss of condition up to 28 weeks is high:

➤ Weigh at least 60 males per pen weekly. Adjust feed according to both condition and bodyweight.

➤ Check feed distribution at least 2 times per week at first and then every week to correct feed allocation and distribution if need be. It is better to attend to the feed distribution to make sure that every male has a normal access to the feeders.


▷ Use a proper grill size 45 mm x 60 mm or adjust pan feeders to control male access to the female feeder. Also ensure that the corners are sealed with covers. Check behaviour to see if males are stealing feed.

▷ Feed males in the morning when the first female feed is being distributed. The feed distribution needs to be uniform all along the feeding system (check if the sensor of the last pan feeder works properly).

▷ The male feeder should be high enough to prevent the females from eating with the males. Consider the risk of under-feeding, e.g. if male activity is very high which will affect energy requirement. The comb will be paler and listless and body conformation will change and be soft. Avoid irreversible moult.



Manual linear feeder for males – not enough stable




Automatic pan feeder for males – too high



Importance of controlling male access to the female feeder. Yellow circle: open feed access for males



House design with female chain feeder and males with manual linear feeder (middle)



Importance of the grill size on female feeders

4. PRODUCTION PERIOD (22-64 WEEKS): HOW TO MAXIMIZE THE PERSISTENCY OF FERTILITY?


OBJECTIVE: MAXIMISE THE FERTILITY RATES UNTIL DEPLETION.

MALE CONDITION/BODYWEIGHT CONTROL

- ▶ Weekly control of male bodyweight (3-5% of the flock and at least 60 birds/pen), male condition (fleshing (shape and amount of breast), footpads, vent, fleshing) is essential.
- ▶ Check the mating activity in the last 2 hours before lights are turned off by sitting on a chair along the sidewalls for at least 30 min. Please seek advice from your Hubbard Technical Manager.
 - ▷ Record the number of mating attempts: it may normally be between 60 and 100 per hour for 1000 females.
 - ▷ Record the number of hens that escape after one mating attempt. It should not be above 30% of the number of mating attempts.

ANALYSIS OF THE WEEKLY RESULTS FOR AN OPTIMAL MALE MANAGEMENT

- ▶ Feed:
 - ▷ Feed should be immediately adjusted at any time that bodyweight drifts away from the recommended objective.
 - ▷ Boost feeding especially when a decrease of the fleshing score is observed e.g. 2 to 3 days with 150% of the current daily feed allocation.
 - ▷ Use of Male 2 diet after 30 weeks of age (lower protein and higher energy content) to meet the weekly growth targets and to allow males to maintain sufficient energy intake without an increased risk of excess fleshing. → See *“Conventional Nutrition Recommendations”*.
 - ▷ Check the physical quality of the feed on a regular basis using the Hubbard feed sieve and the Hubbard software. → See *“Hubbard Feed Sieve Tool”*.
- ▶ Regular culling of poor-quality males.
- ▶ Mating activity could be encouraged by distribution of grain in the late afternoon after 40 weeks of age (e.g. 1g/bird every 2-3 days) and/or adjustment of the sex ratio.
- ▶ **MALE REPLACEMENT:** Biosecurity must be considered before doing this, especially in areas with all health risks. → See Technical bulletin *“Male replacement”*.
 - ▷ Inter-house male replacement does not require importing males from outside. All poor-quality males are taken out and humanely euthanized and replaced by males with similar bodyweight.
 - ▷ External male replacement with young males (25-28 weeks old) involves strict sanitary control of the new males. Between 20 and 30% of the male flock size will be replaced to stimulate the old males and will replace the males with poor physical condition.



Strong and straight legs
Good feet



Successful mating



Red and moist cloaca



Good testicle size

i

[Conventional Nutrition Recommendations](#)

[Hubbard Feed Sieve Tool](#)

[Technical bulletin Male replacement](#)

5. BIOSECURITY

LITTER MANAGEMENT



Litter/bedding storage in an adapted biosecure area to maintain the pathogen-free status (inside the house).



No accumulated litter at the end of the cycle.


SHED TO SHED ACCESS



If corridor between the two houses provide three-zone entry .



If no corridor between the houses, provide one shower unit in each house.



VISITORS



Restrict visitors and adapt quarantine period.



Provide shower facilities.



Provide change of clothes and boots.

VEHICLES



Vehicles must be cleaned prior to entering the farm facility.



DO NOT allow unauthorised vehicles on the farm.



Make sure all feed and gas deliveries stay outside of the fence.



Bags are to be worn on shoes from cars into the shower facilities.

EXTERIOR



Do not treat the concrete or area inside the fence as clean.

INTERIOR



Easy to clean and disinfect concrete floor.



Easy to clean and disinfect equipment. The resting period ideally lasts for at least 10 days.



Ensure good quality water is available.

ANIMAL CONTROL



Keep other livestock away from facility.

PEST CONTROL




Establish regular baiting for rodents and insects, prevent access of wild birds according to local regulations.

KEEP A RECORD



Visitors, feed/litter delivery, cleaning and disinfecting, pest control.



Freezer room: disposal of dead birds in a biosecure freezer.

1 Gas tank

2 Fumigation room

3 Egg storage room

4 Vehicle disinfection unit

5 Staff room

6 Staff room

→ Go forward principle



The performance data contained in this document was obtained from results and experience from our own research flocks and flocks of our customers. In no way does the data contained in this document constitute a warranty or guarantee of the same performance under different conditions of nutrition, density or physical or biological environment. In particular (but without limitation of foregoing) we do not grant any warranties regarding the fitness for purpose, performance, use, nature or quality of the flocks, nor any warranty regarding compliance with local legislation regarding health, welfare, or other aspects of animal production. Hubbard makes no representation as to the accuracy or completeness of the information contained in this document.



AMERICAS
HUBBARD LLC
1070 MAIN STREET
PIKEVILLE, TN 37367 – U.S.A.
TEL. +1 (423) 447 6224
contact.americas@hubbardbreedersusa.com

EUROPE, MIDDLE EAST, AFRICA
HUBBARD S.A.S.
MAUGUÉRAND
22800 LE FOEIL – FRANCE
TEL. +33 2 96 79 63 70
contact.emea@hubbardbreeders.com

ASIA
HUBBARD S.A.S.
MAUGUÉRAND
22800 LE FOEIL – FRANCE
TEL. +33 2 96 79 63 70
contact.asia@hubbardbreeders.com

Hubbard is a registered trademark of Hubbard in the US and other countries.
All other brands and trademarks are the trademarks of their respective owners.

© Hubbard